

Unusual Features of the SARS-CoV-2 Genome Suggesting Sophisticated Laboratory Modification as a Biological Robot Rather than Natural Evolution and Delineation of Its Probable Synthetic Route

Li-Meng Yan¹ and Adrian David Cheok²

¹ University Tokyo

Received: 9 September 2021 Accepted: 1 October 2021 Published: 15 October 2021

Abstract

The COVID-19 pandemic caused by the novel coronavirus SARS-CoV-2 has led to over 4.24 million deaths worldwide and unprecedented decimation of the global economy. Despite its tremendous impact, the origin of SARS-CoV-2 has remained mysterious and controversial. The natural origin theory, although widely accepted, lacks substantial support. The alternative theory that the virus may have come from a research laboratory is, however, strictly censored on peerreviewed scientific journals. Nonetheless, SARS-CoV-2 shows biological characteristics that are inconsistent with a naturally occurring, zoonotic virus. In this report, we describe the genomic, structural, medical, and literature evidence, which, when considered together, strongly contradicts the natural origin theory. The evidence shows that SARS-CoV2 should be a laboratory product created by using bat coronaviruses ZC45 and/or ZXC21 as a template and/or backbone.

Index terms—

1 Introduction

COVID-19 has caused a world-wide pandemic, the scale and severity of which are unprecedented. Despite the tremendous efforts taken by the global community, management and control of this pandemic remains difficult and challenging.

As a coronavirus, SARS-CoV-2 differs significantly from other respiratory and/or zoonotic viruses: it attacks multiple organs; it is capable of undergoing a long period of asymptomatic infection; it is highly transmissible and significantly lethal in high-risk populations; it is well-adapted to humans since the very start of its emergence; it is highly efficient in binding the human ACE2 receptor (hACE2), the affinity of which is greater than that associated with the ACE2 of any other potential host 2,3 .

The origin of SARS-CoV-2 is still the subject of much debate. A widely cited Nature Medicine publication has claimed that SARS-CoV-2 most likely came from nature 4 . However, the article and its central conclusion are now being challenged by scientists from all over the world [5][6][7][8][9][10][11][12][13][14][15] . In addition, authors of this Nature Medicine article show signs of conflict of interests 16,17 , raising further concerns on the credibility of this publication.

The existing scientific publications supporting a natural origin theory rely heavily on a single piece of evidence—a previously discovered bat coronavirus named RaTG13, which shares a 96% nucleotide sequence identity with SARS-CoV-2 18 . However, the existence of RaTG13 in nature and the truthfulness of its reported sequence are being widely questioned [6][7][8][9][19][20][21] . It is noteworthy that scientific journals have clearly censored any dissenting opinions that suggest a nonnatural origin of SARS-CoV-2 8,22 . Because of this censorship, articles

2 A) HAS SARS-COV-2 BEEN SUBJECTED TO IN VITRO MANIPULATION?

questioning either the natural origin of SARS-CoV-2 or the actual existence of RaTG13, although of high quality scientifically, can only exist as preprints [5][6][7][8][9][19][20][21] or other non-peer reviewed articles published on various online platforms [10][11][12][13]23 . Nonetheless, analyses of these reports have repeatedly pointed to severe problems and a probable fraud associated with the reporting of RaTG13 6,8,9, ??921 . Therefore, the theory that fabricated scientific data has been published to mislead the world's efforts in tracing the origin of SARS-CoV-2 has become substantially convincing and is interlocked with the notion that SARS-CoV-2 is of a non-natural origin.

Consistent with this notion, genomic, structural, and literature evidence also suggest a non-natural origin of SARS-CoV-2. In addition, abundant literature C Figure ??: Genomic sequence analysis reveals that bat coronavirus ZC45 is the closest match to SARS-CoV-2. Top: genomic organization of SARS-CoV-2 (2019-nCoV WIV04). Bottom: similarity plot based on the full-length genome of 2019-nCoV WIV04. Full-length genomes of SARS-CoV BJ01, bat SARSr-CoV WIV1, bat SARSr-CoV HKU3-1, bat coronavirus ZC45 were used as reference sequences.

indicates that gain-of-function research has long advanced to the stage where viral genomes can be precisely engineered and manipulated to enable the creation of novel coronaviruses possessing unique properties. In this report, we present such evidence and the associated analyses. Part 1 of the report describes the genomic and structural features of SARS-CoV2, the presence of which could be consistent with the theory that the virus is a product of laboratory modification beyond what could be afforded by simple serial viral passage. Part 2 of the report describes a highly probable pathway for the laboratory creation of SARS-CoV-2, key steps of which are supported by evidence present in the viral genome. Importantly, part 2 should be viewed as a demonstration of how SARS-CoV-2 could be conveniently created in a laboratory in a short period of time using available materials and well-documented techniques. This report is produced by a team of experienced scientists using our combined expertise in virology, molecular biology, structural biology, computational biology, vaccine development, and medicine.

2 a) Has SARS-CoV-2 been subjected to in vitro manipulation?

We present three lines of evidence to support our contention that laboratory manipulation is part of the history of SARS-CoV-2:

i. The genomic sequence of SARS-CoV-2 is suspiciously similar to that of a bat coronavirus discovered by military laboratories in the Third Military Medical University (Chongqing, China) and the Research Institute for Medicine of Nanjing Command (Nanjing, China). ii. The receptor-binding motif (RBM) within the Spike protein of SARS-CoV-2, which determines the host specificity of the virus, resembles that of SARS-CoV from the 2003 epidemic in a suspicious manner. Genomic evidence suggests that the RBM has been genetically manipulated. iii. SARS-CoV-2 contains a unique furin-cleavage site in its Spike protein, which is known to greatly enhance viral infectivity and cell tropism. Yet, this cleavage site is completely absent in this particular class of coronaviruses found in nature. In addition, rare codons associated with this additional sequence suggest the strong possibility that this furin-cleavage site is not the product of natural evolution and could have been inserted into the SARS-CoV-2 genome artificially by techniques other than simple serial passage or multi-strain recombination events inside co-infected tissue cultures or animals.

i. Genomic sequence analysis reveals that ZC45, or a closely related bat coronavirus, should be the backbone used for the creation of SARS-CoV-2

The structure of the ~30,000 nucleotides-long SARS-CoV-2 genome is shown in Figure ?. Searching the NCBI sequence database reveals that, among all known coronaviruses, there were two related bat coronaviruses, ZC45 and ZXC21, that share the highest sequence identity with SARS-CoV-2 (each bat coronavirus is ~89% identical to SARS-CoV-2 on the nucleotide level). Similarity between the genome of SARS-CoV-2 and those of representative ? coronaviruses is depicted in Figure ?. ZXC21, which is 97% identical to and shares a very similar profile with ZC45, is not shown. Note that the RaTG13 virus is excluded from this analysis given the strong evidence suggesting that its sequence may have been fabricated and the virus does not exist in nature 2,[6][7][8][9] .

Orf8 is an accessory protein, the function of which is largely unknown in most coronaviruses, although recent data suggests that Orf8 of SARS-CoV-2 mediates the evasion of host adaptive immunity by down regulating MHC-I 24 . Normally, Orf8 is poorly conserved in coronaviruses 25 . Sequence blast indicates that, while the Orf8 proteins of ZC45/ZXC21 share a 94.2% identity with SARS-CoV-2 Orf8, no other coronaviruses share more than 58% identity with SARS-CoV-2 on this particular protein. The very high homology here on the normally poorly conserved Orf8 protein is highly unusual. The coronavirus E protein is a structural protein, which is embedded in and lines the interior of the membrane envelope of the virion 26 . The E protein is tolerant of mutations as evidenced in both SARS (Figure 2A) and related bat coronaviruses (Figure 2B). This tolerance to amino acid mutations of the E protein is further evidenced in the current SARS-CoV-2 pandemic. After only a short two-month spread of the virus since its outbreak in humans, the E proteins in SARS-CoV-2 have already undergone mutational changes. Sequence data obtained during the month of April reveals that mutations have occurred at four different locations in different strains (Figure 2C). Consistent with this finding, sequence blast analysis indicates that, with the exception of SARS-CoV-2, no known coronaviruses share 100% amino acid sequence identity on the E protein with ZC45/ZXC21 (suspicious coronaviruses published after the start of

104 the current pandemic are excluded 18,[27][28][29][30][31]). Although 100% identity on the E protein has been
105 observed between SARS-CoV and certain SARS-related bat coronaviruses, none of those pairs simultaneously
106 share over 83% identity on the Orf8 protein 32 . Therefore, the 94.2% identity on the Orf8 protein, 100% identity
107 on the E protein, and the overall genomic/amino acid-level resemblance between SARS-CoV-2 and ZC45/ZXC21
108 are highly unusual. Such When SARS-CoV-2 and ZC45/ZXC21 are compared on the amino acid level, a high
109 sequence identity is observed for most of the proteins. The Nucleocapsid protein is 94% identical. The Membrane
110 protein is 98.6% identical. The S2 portion (2nd half) of the Spike protein is 95% identical. Importantly, the
111 Orf8protein is 94.2% identical and the E protein is 100% identical.

112 evidence, when considered together, is consistent with a hypothesis that the SARS-CoV-2 genome has an origin
113 based on the use of ZC45/ZXC21 as a backbone and/or template for genetic gain-of-function modifications.

114 Importantly, ZC45 and ZXC21 are bat corona viruses that were discovered (between July 2015 and February
115 2017), isolated, and characterized by military research laboratories in the Third Military MedicalUniversity
116 (Chongqing, China) and the Research Institute for Medicine of Nanjing Command (Nanjing, China). The data
117 and associated work were published in 2018 33,34 . Clearly, this backbone/template, which is essential for the
118 creation of SARS-CoV-2, exists in these and other related research laboratories.

119 What strengthens our contention further is the published RaTG13 virus 18 , the genomic sequence of which is
120 reportedly 96% identical to that of SARS-CoV-2. While suggesting a natural origin of SARSCoV-2, the RaTG13
121 virus also diverted the attention of both the scientific field and the general public away from ZC45/ZXC21
122 4,18 . In fact, a Chinese BSL-3 lab (the Shanghai Public Health Clinical Centre), which published a Nature
123 article reporting a conflicting close phylogenetic relationship between SARSCoV-2 and ZC45/ZXC21 rather than
124 with RaTG13 35 , was quickly shut down for "rectification" 36 . It is believed that the researchers of that
125 laboratory were being punished for having disclosed the SARS-CoV2-ZC45/ZXC21 connection. On the other
126 hand, substantial evidence has accumulated, pointing to severe problems associated with the reported sequence
127 of RaTG13 as well as questioning the actual existence of this bat virus in nature 6,7,[19][20][21] . A very recent
128 publication also indicated that the receptor-binding domain (RBD) of the RaTG13's Spike protein could not bind
129 ACE2 of two different types of horseshoe bats (they closely relate to the horseshoe bat *R. affinis*, RaTG13's alleged
130 natural host) 2 , implicating the inability of RaTG13 to infect horseshoe bats. This finding further substantiates
131 the suspicion that the reported sequence of RaTG13 could have been fabricated as the Spike protein encoded by
132 this sequence does not seem to carry the claimed function. The fact that a virus has been fabricated to shift the
133 attention away from ZC45/ZXC21 speaks for an actual role of ZC45/ZXC21 in the creation of SARS-CoV-2.

134 ii. The receptor-binding motif of SARS-CoV-2 Spike cannot be born from nature and should have been created
135 through genetic engineering The Spike proteins decorate the exterior of the coronavirus particles. They play an
136 important role in infection as they mediate the interaction with host cell receptors and thereby help determine
137 the host range and tissue tropism of the virus. The Spike protein is split receptor. In both SARS-CoV and SARS-
138 CoV-2 infections, the host cell receptor is hACE2. Within S1, a segment of around 70 amino acids makes direct
139 contacts with hACE2 and is correspondingly named the receptorbinding motif (RBM) (Figure 3C). In SARS-
140 CoV and SARS-CoV-2, the RBM fully determines the interaction with hACE2. The C-terminal half of the
141 Spike protein is named S2. The main function of S2 includes maintaining trimer formation and, upon successive
142 protease cleavages at the S1/S2 junction and a downstream S2' position, mediating membrane fusion to enable
143 cellular entry of the virus. Similar to what is observed for other viral proteins, S2 of SARS-CoV-2 shares a high
144 sequence identity (95%) with S2 of ZC45/ZXC21. In stark contrast, between SARS-CoV-2 and ZC45/ZXC21,
145 the S1 protein, which dictates which host (human or bat) the virus can infect, is much less conserved with the
146 amino acid sequence identity being only 69%.

147 Figure 4 shows the sequence alignment of the Spike proteins from six ? coronaviruses. Two are viruses
148 isolated from the current pandemic (Wuhan-Hu-1, 2019-nCoV_USA-AZ1); two are the suspected template
149 viruses (Bat_CoV_ZC45, Bat_CoV_ZXC21); two are SARS coronaviruses (SARS_GZ02, SARS). The RBM is
150 highlighted in between two orange lines. Clearly, despite the high sequence identity for the overall genomes, the
151 RBM of SARS-CoV-2 differs significantly from those of ZC45 and ZXC21. Intriguingly, the RBM of SARS-CoV-2
152 resembles, on a great deal, the RBM of SARS Spike. Although this is not an exact "copy and paste", careful
153 examination of the Spike-hACE2 structures 37,38 reveals that all residues essential for either hACE2 binding or
154 protein folding (orange sticks in Figure 3C and what is highlighted by red short lines in Figure 4) are "kept". Most
155 of these essential residues are precisely preserved, including those involved in disulfide bond formation (C467,
156 C474) and electrostatic interactions (R444, E452, R453, D454), which are pivotal for the structural integrity of
157 the RBM (Figure 3C and 4). The few changes within the group of essential residues are As elaborated below, the
158 way that SARS-CoV-2 RBM resembles SARS-CoV RBM and the overall sequence conservation pattern between
159 SARS-CoV-2 and ZC45/ZXC21 are highly unusual. Collectively, this suggests that portions of the SARS-CoV-2
160 genome have not been derived from natural quasi-species viral particle evolution.

161 If SARS-CoV-2 does indeed come from natural evolution, its RBM could have only been acquired in one of
162 the two possible routes: 1) an ancient recombination event followed by convergent evolution or 2) a natural
163 recombination event that occurred fairly recently.

164 In the first scenario, the ancestor of SARS-CoV-2, a ZC45/ZXC21-like bat coronavirus would have recombined
165 and "swapped" its RBM with a coronavirus carrying a relatively "complete" RBM (in reference to SARS).
166 This recombination would result in a novel ZC45/ZXC21-like coronavirus with all the gaps in its RBM "filled"

2 A) HAS SARS-COV-2 BEEN SUBJECTED TO IN VITRO MANIPULATION?

167 (Figure 4). Subsequently, the virus would have to adapt extensively in its new host, where the ACE2 protein is
168 highly homologous to hACE2. Random mutations across the genome would have to have occurred to eventually
169 shape the RBM to its current form -resembling SARS-CoV RBM in a highly intelligent manner. However, this
170 convergent evolution process would also result in the accumulation of a large amount of mutations in other parts
171 of the genome, rendering the overall sequence identity relatively low. The high sequence identity between SARS-
172 CoV-2 and ZC45/ ZXC21 on various proteins (94-100% identity) do not support this scenario and, therefore,
173 clearly indicates that SARS-CoV2 carrying such an RBM cannot come from a ZC45/ZXC21-like bat coronavirus
174 through this convergent evolutionary route.

175 In the second scenario, the ZC45/ZXC21-like coronavirus would have to have recently recombined and swapped
176 its RBM with another coronavirus that had successfully adapted to bind an animal ACE2 highly homologous to
177 hACE2. The likelihood of such an event depends, in part, on the general requirements of natural recombination:
178 1) that the two different viruses share significant sequence similarity; 2) that they must coinfect and be present
179 in the same cell of the same animal; 3) that the recombinant virus would not be cleared by the host or make the
180 host extinct; 4) that the recombinant virus eventually would have to become stable and transmissible within the
181 host species.

182 In regard to this recent recombination scenario, the animal reservoir could not be bats because the ACE2
183 proteins in bats are not homologous enough to hACE2 and therefore the adaptation would not be able to yield an
184 RBM sequence as seen in SARS-CoV-2. This animal reservoir also could not be humans as the ZC45/ZXC21-like
185 coronavirus would not be able to infect humans. In addition, there has been no evidence of any SARS-CoV-2
186 or SARS-CoV-2-like virus circulating in the human population prior to late 2019. Intriguingly, according to a
187 recent bioinformatics study, SARS-CoV-2 was well-adapted for humans since the start of the outbreak 1 .

188 Only one other possibility of natural evolution remains, which is that the ZC45/ZXC21-like virus and a
189 coronavirus containing a SARS-like RBM could have recombined in an intermediate host where the ACE2 protein
190 is homologous to hACE2. Several laboratories have reported that some of the Sunda pangolins smuggled into
191 China from Malaysia carried coronaviruses, the receptor-binding domain (RBD) of which is almost identical to
192 that of SARS-CoV-2 [27][28][29]31 . They then went on to suggest that pangolins are the likely intermediate
193 host for SARS-CoV-2 [27][28][29]31 . However, recent independent reports have found significant flaws() G Year
194 2021

195 Unusual Features of the SARS-CoV-2 Genome Suggesting Sophisticated Laboratory Modification as a
196 Biological Robot Rather than Natural Evolution and Delineation of Its Probable Synthetic Route in this data
197 [40][41][42] . Furthermore, contrary to these almost exclusively hydrophobic "substitutions" (I428?L, L443?F,
198 F460?Y, L472?F, Y484?Q), which should not affect either protein folding or the hACE2-interaction. At the same
199 time, majority of the amino acid residues that are non-essential have "mutated" (Figure 4, RBM residues not
200 labeled with short red lines). Judging from this sequence analysis alone, we were convinced early on that not
201 only would the SARS-CoV-2 Spike protein bind hACE2 but also the binding would resemble, precisely, that
202 between the original SARS Spike protein and hACE2 23 . Recent structural work has confirmed our prediction
203 39 . reports [27][28][29]31 , no coronaviruses have been detected in Sunda pangolin samples collected for over a
204 decade in Malaysia and Sabah between 2009 and 2019 43 . A recent study also showed that the RBD, which is
205 shared between SARS-CoV-2 and the reported pangolin coronaviruses, binds to hACE2 ten times stronger than
206 to the pangolin ACE2 2 , further dismissing pangolins as the possible intermediate host. Finally, an in silico
207 study, while echoing the notion that pangolins are not likely an intermediate host, also indicated that none of the
208 animal ACE2 proteins examined in their study exhibited more favorable binding potential to the SARS-CoV-2
209 Spike protein than hACE2 did 3 . This last study virtually exempted all animals from their suspected roles
210 as an intermediate host 3 , which is consistent with the observation that SARS-CoV-2 was well-adapted for
211 humans from the start of the outbreak 1 . This is significant because these findings collectively suggest that
212 no intermediate host seems to exist for SARS-CoV-2, which at the very least diminishes the possibility of a
213 recombinant event occurring in an intermediate host.

214 Even if we ignore the above evidence that no proper host exists for the recombination to take place and
215 instead assume that such a host does exist, it is still highly unlikely that such a recombination event could occur
216 in nature.

217 As we have described above, if natural recombination event is responsible for the appearance of SARSCoV-
218 2, then the ZC45/ZXC21-like virus and a coronavirus containing a SARS-like RBM would have to recombine
219 in the same cell by swapping the S1/RBM, which is a rare form of recombination. Furthermore, since SARS
220 has occurred only once in human history, it would be at least equally rare for nature to produce a virus that
221 resembles SARS in such an intelligent manner -having an RBM that differs from the SARS RBM only at a few
222 non-essential sites (Figure 4). The possibility that this unique SARS-like coronavirus would reside in the same cell
223 with the ZC45/ZXC21-like ancestor virus and the two viruses would recombine in the "RBMSwapping" fashion
224 is extremely low. Importantly, this, and the other recombination event described below in section 1.3 (even more
225 impossible to occur in nature), would both have to happen to produce a Spike as seen in SARS-CoV-2.

226 While the above evidence and analyses together appear to disapprove a natural origin of SARS-CoV2's RBM,
227 abundant literature shows that gain-of-function research, where the Spike protein of a coronavirus was specifically
228 engineered, has repeatedly led to the successful generation of humaninfecting coronaviruses from coronaviruses
229 of non-human origin [44][45][46][47] .

Record also shows that research laboratories, for example, the Wuhan Institute of Virology (WIV), have successfully carried out such studies working with US researchers 45 and also working alone 47 . In addition, the WIV has engaged in decades-long coronavirus surveillance studies and therefore owns the world’s largest collection of coronaviruses. Evidently, the technical barrier is non-existent for the WIV and other related laboratories to carry out and succeed in such Spike/RBM engineering and gain-of function research. 39 . The RBM highlighted in blue (bottom) is from the Spike protein (RBM: 424-494) of SARS-BJ01 (AY278488.2), which was swapped by the Shi lab into the Spike proteins of different bat coronaviruses replacing the corresponding segments 47 .

Strikingly, consistent with the RBM engineering theory, we have identified two unique restriction sites, EcoRI and BstEII, at either end of the RBM of the SARS-CoV-2 genome, respectively (Figure 5A). These two sites, which are popular choices of everyday molecular cloning, do not exist in the rest of this spikegene. This particular setting makes it extremely convenient to swap the RBM within spike, providing a quick way to test different RBMs and the corresponding Spike proteins.

Such EcoRI and BstEII sites do not exist in the spike genes of other ? coronaviruses, which strongly indicates that they were unnatural and were specifically introduced into this spike gene of SARS-CoV-2 for the convenience of manipulating the critical RBM. Although ZC45 spike also does not have these two sites (Figure 5B), they can be introduced very easily as described in part 2 of this report.

It is noteworthy that introduction of the EcoRI site here would change the corresponding amino acids from -WNT-to -WNS-(Figure 5AB). As far as we know, all SARS and SARS-like bat coronaviruses exclusively carry a T (threonine) residue at this location. SARS-CoV-2 is the only exception in that this T has mutated to an S (serine), save the suspicious RaTG13 and pangolin coronaviruses published after the outbreak 48 .

Once the restriction sites were successfully introduced, the RBM segment could be swapped conveniently using routine restriction enzyme digestion and ligation. Although alternative cloning techniques may leave no trace of genetic manipulation (Gibson assembly as one example), this oldfashioned approach could be chosen because it offers a great level of convenience in swapping this critical RBM.

Given that RBM fully dictates hACE2-binding and that the SARS RBM-hACE2 binding was fully characterized by high-resolution structures (Figure 3) 37,38 , this RBM-only swap would not be any riskier than the full Spike swap. In fact, the feasibility of this RBMswap strategy has been proven 39,47 . In 2008, Dr. Zhengli Shi’s group swapped a SARS RBM into the Spike proteins of several SARS-like bat coronaviruses after introducing a restriction site into a codon-optimized spike gene (Figure 5C) 47 . They then validated the binding of the resulted chimeric Spike proteins with hACE2. Furthermore, in a recent publication, the RBM of SARS-CoV-2 was swapped into the receptor-binding domain (RBD) of SARSCoV, resulting in a chimeric RBD fully functional in binding hACE2 (Figure 5C) 39 . Strikingly, in bothcases, the manipulated RBM segments resemble almost exactly the RBM defined by the positions of theEcoRI and BstEII sites (Figure 5C). Although cloning details are lacking in both publications 39,47 , it is conceivable that the actual restriction sites may vary depending on the spike gene receiving the RBM insertion as well as the convenience in introducing unique restriction site(s) in regions of interest. It is noteworthy that the corresponding author of this recent publication 39 , Dr. Fang Li, has been an active collaborator of Dr. Zhengli Shi since 2010 [49][50][51][52][53] . Dr. Li was the first person in the world to have structurally elucidated the binding between SARS-CoV RBD and hACE2 38 and has been the leading expert in the structural understanding of Spike-ACE2 interactions 38,39,[53][54][55][56] . The striking finding of EcoRI and BstEII restriction sites at either end of the SARS-CoV-2 RBM, respectively, and the fact that the same RBM region has been swapped both by Dr. Shi and by her long-term collaborator, respectively, using restriction enzyme digestion methods are unlikely a coincidence. Rather, it is the smoking gun proving that theRBM/Spike of SARS-CoV-2 is a product of genetic manipulation.

Although it may be convenient to copy the exact sequence of SARS RBM, it would be too clear a sign of artificial design and manipulation. The more deceiving approach would be to change a few nonessential residues, while preserving the ones critical for binding. This design could be well-guided by the high-resolution structures (Figure 3) 37,38 . This way, when the overall sequence of the RBM would appear to be more distinct from that of the SARS RBM, the hACE2-binding ability would be well-preserved. We believe that all of the crucial residues (residues labeled with red sticks in Figure 4, which are the same residues shown in sticks in Figure 3C) should have been ”kept”. As described earlier, while some should be direct preservation, some should have been switched to residues with similar properties, which would not disrupt hACE2-binding and may even strengthen the association further. Importantly, changes might have been made intentionally at non-essential sites, making it less like a ”copy and paste” of the SARS RBM.

iii. An unusual furin-cleavage site is present in the Spike protein of SARS-CoV-2 and is associated with the augmented virulence of the virus Another unique motif in the Spike protein of SARS-CoV-2 is a polybasic furin-cleavage site located at the S1/S2 junction (Figure 4, segment in between two green lines). Such a site can be recognized and cleaved by the furin protease. Within the lineage B of ? coronaviruses and with the exception of SARSCoV-2, no viruses contain a furin-cleavage site at the S1/S2 junction (Figure 6) 57 . In contrast, furincleavage site at this location has been observed in other groups of coronaviruses 57,58 . Certain selective pressure seems to be in place that prevents the lineage B of ? coronaviruses from acquiring or maintaining such a site in nature. As previously described, during the cell entry process, the Spike protein is first cleaved at the S1/S2 junction. This step, and a subsequent cleavage downstream that exposes the fusion peptide, are both mediated by host proteases. The presence or absence of these proteases in different cell types greatly affects the

3 B) DELINEATION OF A SYNTHETIC ROUTE OF SARS-COV-2

293 cell tropism and presumably the pathogenicity of the viral infection. Unlike other proteases, furin protease is
294 widely expressed in many types of cells and is present at multiple cellular and extracellular locations. Importantly,
295 the introduction of a furin-cleavage site at the S1/S2 junction could significantly enhance the infectivity of a virus
296 as well as greatly expand its cell tropism -a phenomenon well-documented in both influenza viruses and other
297 coronaviruses [59][60][61][62][63][64][65] .

298 If we leave aside the fact that no furin-cleavage site is found in any lineage B ? coronavirus in nature and
299 instead assume that this site in SARS-CoV-2 is a result of natural evolution, then only one evolutionary pathway
300 is possible, which is that the furin-cleavage site has to be derived from a homologous recombination event.
301 Specifically, an ancestor ? coronavirus containing no furin-cleavage site would have to recombine with a closely
302 related coronavirus that does contain a furin cleavage site.

303 However, two facts disfavor this possibility. First, although some coronaviruses from other groups or lineages do
304 contain polybasic furin-cleavage sites, none of them contains the exact polybasic sequence present in SARS-CoV-
305 2 (-PRRAR/SVA-). Second, between SARS-CoV-2 and any coronavirus containing a legitimate furin cleavage
306 site, the sequence identity on Spike is no more than 40% 66 . Such a low level of sequence identity rules out
307 the possibility of a successful homologous recombination ever occurring between the ancestors of these viruses.
308 Therefore, the furin-cleavage site within the SARS-CoV-2 Spike protein is unlikely to be of natural origin and
309 instead should be a result of laboratory modification.

310 Consistent with this claim, a close examination of the nucleotide sequence of the furin-cleavage site in SARS-
311 CoV-2 spike has revealed that the two consecutive Arg residues within the inserted sequence (PRR Figure ??:
312 Two consecutive Arg residues in the -PRRA-insertion at the S1/S2 junction of SARS-CoV-2 Spike are both
313 coded by a rare codon, CGG. A FauI restriction site, 5'-(N) 6 GCGGG-3', is embedded in the coding sequence
314 of the "inserted" PRRA segment, which may be used as a marker to monitor the preservation of the introduced
315 furin cleavage site.

316 A-) are both coded by the rare codon CGG (least used codon for Arg in SARS-CoV-2) (Figure ??) 8 . In
317 fact, this CGGCGG arrangement is the only instance found in the SARS-CoV-2 genome where this rare codon
318 is used in tandem. This observation strongly suggests that this furin-cleavage site should be a result of genetic
319 engineering. Adding to the suspicion, a FauI restriction site is formulated by the codon choices here, suggesting
320 the possibility that the restriction fragment length polymorphism, a technique that a WIV lab is proficient at
321 67 , could have been involved. There, the fragmentation pattern resulted from FauI digestion could be used to
322 monitor the preservation of the furin cleavage site in Spike as this furin cleavage site is prone to deletions in vitro
323 68,69 . Specifically, RT-PCR on the spike gene of the recovered viruses from cell cultures or laboratory animals
324 could be carried out, the product of which would be subjected to FauI digestion. Viruses retaining or losing the
325 furin-cleavage site would then yield distinct patterns, allowing convenient tracking of the virus(es) of interest.

326 In addition, although no known coronaviruses contain the exact sequence of -PRRAR/SVA-that is present in
327 the SARS-CoV-2 Spike protein, a similar -RRAR/AR-sequence has been observed at the S1/S2 junction of the
328 Spike protein in a rodent coronavirus, AcCoV-JC34, which was published by Dr. Zhengli Shi in 2017 70 . It
329 is evident that the legitimacy of -RRAR-as a functional furin-cleavage site has been known to the WIV experts
330 since 2017.

331 The evidence collectively suggests that the furin-cleavage site in the SARS-CoV-2 Spike protein may not have
332 come from nature and could be the result of genetic manipulation. The purpose of this manipulation could have
333 been to assess any potential enhancement of the infectivity and pathogenicity of the laboratory made coronavirus
334 [59][60][61][62][63][64] . Indeed, recent studies have confirmed that the furin-cleavage site does confer significant
335 pathogenic advantages to SARS-CoV-2 57,68 .

336 iv. Discussion Evidence presented in this part reveals that certain aspects of the SARS-CoV-2 genome are
337 extremely difficult to reconcile to being a result of natural evolution. The alternative theory we suggest is that the
338 virus may have been created by using ZC45/ZXC21 bat coronavirus(es) as the backbone and/or template. The
339 Spike protein, especially the RBM within it, should have been artificially manipulated, upon which the virus has
340 acquired the ability to bind hACE2 and infect humans. This is supported by the finding of a unique restriction
341 enzyme digestion site at either end of the RBM. An unusual furin-cleavage site may have been introduced and
342 inserted at the S1/S2 junction of the Spike protein, which contributes to the increased virulence and pathogenicity
343 of the virus. These transformations have then staged the SARSCoV-2 virus to eventually become a highly-
344 transmissible, onset-hidden, lethal, sequelae unclear, and massively disruptive pathogen.

345 Evidently, the possibility that SARS-CoV-2 could have been created through gain-of-function manipulations
346 at the WIV is significant and should be investigated thoroughly and independently.

3 b) Delineation of a synthetic route of SARS-CoV-2

348 In the second part of this report, we describe a synthetic route of creating SARS-CoV-2 in a laboratory setting.
349 It is postulated based on substantial literature support as well as genetic evidence present in the SARS-CoV-2
350 genome. Although steps presented herein should not be viewed as exactly those taken, we believe that key
351 processes should not be much different. Importantly, our work here should serve as a demonstration of how
352 SARS-CoV-2 can be designed and created conveniently in research laboratories by following proven concepts and
353 using well-established techniques.

354 Importantly, research labs, both in Hong Kong and in mainland China, are leading the world in coronavirus

355 research, both in terms of resources and on the research outputs. The latter is evidenced not only by the
356 large number of publications that they have produced over the past two decades but also by their milestone
357 achievements in the field: they were the first to identify civets as the intermediate host for SARS-CoV and
358 isolated the first strain of the virus 71 ; they were the first to uncover that SARS-CoV originated from bats 72,73
359 ; they revealed for the first time the antibody-dependent enhancement (ADE) of SARS-CoV infections 74 ; they
360 have contributed significantly in understanding MERS in all domains (zoonosis, virology, and clinical studies)
361 [75][76][77][78][79] ; they made several breakthroughs in SARS-CoV-2 research 18,35,80 . Last but not least,
362 they have the world's largest collection of coronaviruses (genomic sequences and live viruses). The knowledge,
363 expertise, and resources are all readily available within the Hong Kong and mainland research laboratories (they
364 collaborate extensively) to carry out and accomplish the work described below. In this sub-section, we outline
365 the possible overall strategy and major considerations that may have been formulated at the designing stage of
366 the project.

367 Once they have chosen a template virus, they would first need to engineer, through molecular cloning, the
368 Spike protein so that it can bind hACE2. The concept and cloning techniques involved in this manipulation have
369 been well-documented in the literature [44][45][46]84,86 . With almost no risk of failing, the template bat virus
370 could then be converted to a coronavirus that can bind hACE2 and infect humans [44][45][46] .

371 Second, they would use molecular cloning to introduce a furin-cleavage site at the S1/S2 junction of Spike.
372 This manipulation, based on known knowledge 60,61,65 , would likely produce a strain of coronavirus that is a
373 more infectious and pathogenic.

374 Third, they would produce an ORF1b gene construct. The ORF1b gene encodes the polyprotein Orf1b, which
375 is processed post-translationally to produce individual viral proteins: RNA-dependent RNA polymerase (RdRp),
376 helicase, guanidine-N7 methyl transferase, uridylate-specific endoribonuclease, and 2'-O-methyltransferase. All
377 of these proteins are parts of the replication machinery of the virus. Among them, the RdRp protein is the
378 most crucial one and is highly conserved among coronaviruses. Importantly, Dr. Zhengli Shi's laboratory uses a
379 PCR protocol, which amplifies a particular fragment of the RdRp gene, as their primary method to detect the
380 presence of coronaviruses in raw samples (bat fecal swap, feces, etc). As a result of this practice, the Shi group
381 has documented the sequence information of this short segment of RdRp for all coronaviruses that they have
382 successfully detected and/or collected.

383 Here, the genetic manipulation is less demanding or complicated because Orf1b is conserved and likely Orf1b
384 from any ? coronavirus would be competent enough to do the work. However, we believe that they would want
385 to introduce a particular Orf1b into the virus for one of the two possible reasons:

386 1. Since many phylogenetic analyses categorize coronaviruses based on the sequence similarity of the RdRp
387 gene only 18,31,35,83,87 , having a different RdRp in the genome therefore could ensure that SARS-CoV-2 and
388 ZC45/ZXC21 are separated into different groups/sub-lineages in phylogenetic studies. Choosing an RdRp gene,
389 however, is convenient because the short RdRp segment sequence has been recorded for all coronaviruses ever
390 collected/detected. Their final choice was the RdRp sequence from bat coronavirus RaBtCoV/4991, which was
391 discovered in 2013. For RaBtCoV/4991, the only information ever published was the sequence of its short RdRp
392 segment 83 , while neither its full genomic sequence nor virus isolation were ever reported. After amplifying the
393 RdRp segment (or the whole ORF1b gene) of RaBatCoV/4991, they would have then used it for subsequent
394 assembly and creation of the genome of SARS-CoV-2. Small changes in the RdRp sequence could either be
395 introduced at the beginning (through DNA synthesis) or be generated viapassages later on. On a separate
396 track, when they were engaged in the fabrication of the RaTG13 sequence, they could have started with the
397 short RdRp segment of RaBtCoV/4991 without introducing any changes to its sequence, resulting in a 100%
398 nucleotide sequence identity between the two viruses on this short RdRp segment 83 . This RaTG13 virus could
399 then be claimed to have been discovered back in 2013. 2. The RdRp protein from RaBatCoV/4991 is unique in
400 that it is superior than RdRp from any other ? coronavirus for developing antiviral drugs. RdRp has no homologs
401 in human cells, which makes this essential viral enzyme a highly desirable target for antiviral development. As
402 an example, Remdesivir, which is currently undergoing clinical trials, targets RdRp. When creating a novel and
403 human-targeting virus, they would be interested in developing the antidote as well. Even though drug discovery
404 like this may not be easily achieved, it is reasonable for them to intentionally incorporate a RdRp that is more
405 amenable for antiviral drug development.

406 Fourth, they would use reverse genetics to assemble the gene fragments of spike, ORF1b, and the rest of the
407 template ZC45 into a cDNA version of the viral genome. They would then carry out in vitro transcription to
408 obtain the viral RNA genome. Transfection of the RNA genome into cells would allow the recovery of live and
409 infectious viruses with the desired artificial genome.

410 To engineer and create a human-targeting coronavirus, they would have to pick a bat coronavirus as the
411 template/backbone. This can be conveniently done because many research labs have been actively collecting
412 bat coronaviruses over the past two decades 32,33,70,72,[81][82][83][84][85] . However, this template virus ideally
413 should not be one from Dr. Zhengli Shi's collections, considering that she is widely known to have been engaged
414 in gain-of-function studies on coronaviruses. Therefore, ZC45 and/or ZXC21, novel bat coronaviruses discovered
415 and owned by military laboratories 33 , would be suitable as the template/backbone. It is also possible that
416 these military laboratories had discovered other closely related viruses from the same location and kept some

417 unpublished. Therefore, the actual template could be ZC45, or ZXC21, or a close relative of them. The postulated
418 pathway described below would be the same regardless of which one of the three was the actual template.

419 Fifth, they would carry out characterization and optimization of the virus strain(s) to improve the fitness,
420 infectivity, and overall adaptation using serial passage in vivo. One or several viral strains that meet certain
421 criteria would then be obtained as the final product(s).

422 4 d) A postulated synthetic route for the creation of SARS- 423 CoV-2

424 In this sub-section, we describe in more details how each step could be carried out in a laboratory setting using
425 available materials and routine molecular, cellular, and virologic techniques. A diagram of this process is shown
426 in Figure 8. We estimate that the whole process could be completed in approximately 6 months.

427 Step 1: Engineering the RBM of the Spike for hACE2binding (1.5

428 5 months)

429 The Spike protein of a bat coronavirus is either incapable of or inefficient in binding hACE2 due to the missing
430 of important residues within its RBM. This can be exemplified by the RBM of the template virus ZC45 (Figure
431 4). The first and most critical step in the creation of SARS-CoV-2 is to engineer the Spike so that it acquires
432 the ability to bind hACE2. As evidenced in the literature, such manipulations have been carried out repeatedly
433 in research laboratories since 2008 44 , which successfully yielded engineered coronaviruses with the ability to
434 infect human cells [44][45][46]88,89 . Although there are many possible ways that one can engineer the Spike
435 protein, we believe that what was actually undertaken was that they replaced the original RBM with a designed
436 and possibly optimized RBM using SARS' RBM as a guide. As described in part 1, this theory is supported
437 by our observation that two unique restriction sites, EcoRI and BstEII, exist at either end of the RBM in the
438 SARS-CoV-2 genome (Figure 5A) and by the fact that such RBM-swap has been successfully carried out by Dr.
439 Zhengli Shi and by her long-term collaborator and structure biology expert, Dr. Fang Li 39,47 .

440 Although ZC45 spike does not contain these two restriction sites (Figure 5B), they can be introduced very
441 easily. The original spike gene would be either amplified with RT-PCR or obtained through DNA synthesis (some
442 changes could be safely introduced to certain variable regions of the sequence) followed by PCR. The gene would
443 then be cloned into a plasmid using restriction sites other than EcoRI and BstEII.

444 Once in the plasmid, the spike gene can be modified easily. First, an EcoRI site can be introduced by converting
445 the highlighted "gaacac" sequence (Figure 5B) to the desired "gaattc" (Figure 5A). The difference between them
446 are two consecutive nucleotides. Using the commercially available Quik Change Site-Directed Mutagenesis kit,
447 such a dinucleotide mutation can be generated in no more than one week. Subsequently, the BstEII site could
448 be similarly introduced at the other end of the RBM. Specifically, the "gaatacc" sequence (Figure 5B) would be
449 converted to the desired "gggtacc" (Figure 5A), which would similarly require a week of time.

450 Once these restriction sites, which are unique within the spike gene of SARS-CoV-2, were successfully
451 introduced, different RBM segments could be swapped in conveniently and the resulting Spike protein
452 subsequently evaluated using established assays.

453 As described in part 1, the design of an RBM segment could be well-guided by the high-resolution structures
454 (Figure 3) 37,38 , yielding a sequence that resembles the SARS RBM in an intelligent manner. When carrying out
455 the structure-guided design of the RBM, they would have followed the routine and generated a few (for example
456 a dozen) such RBMs with the hope that some specific variant(s) may be superior than others in binding hACE2.
457 Once the design was finished, they could have each of the designed RBM genes commercially synthesized (quick
458 and very affordable) with an EcoRI site at the 5'-end and a BstEII site at the 3'-end. These novel RBM genes
459 could then be cloned into the spike gene, respectively. The gene synthesis and subsequent cloning, which could
460 be done in a batch mode for the small library of designed RBMs, would take approximately one month.

461 These engineered Spike proteins might then be tested for hACE2-binding using the established pseudotype
462 virus infection assays 45,49,50 . The engineered Spike with good to exceptional binding affinities would be
463 selected. (Although not necessary, directed evolution could be involved here (error-prone PCR on the RBM
464 gene), coupled with either an in vitro binding assay 39,90 or a pseudotype virus infection assay 45,49,50 , to
465 obtain an RBM that binds hACE2 with exceptional affinity.)

466 Given the abundance of literature on Spike engineering [44][45][46]84,86 and the available high-resolution
467 structures of the Spike-hACE2 complex 37,38 , the success of this step would be very much guaranteed. By the
468 end of this step, as desired, a novel spike gene would be obtained, which encodes a novel Spike protein capable
469 of binding hACE2 with high affinity.

470 Step 2: Engineering a furin-cleavage site at the S1/S2 junction (0.5 month)

471 The product from Step 1, a plasmid containing the engineered spike, would be further modified to include a
472 furin-cleavage site (segment indicated by green lines in Figure 4) at the S1/S2 junction. This short stretch of
473 gene sequence can be conveniently inserted using several routine cloning techniques, including Quik Change Site-
474 Directed PCR 60 , overlap PCR followed by restriction enzyme digestion and ligation 91 , or Gibson assembly.
475 None of these techniques would leave any trace in the sequence. Whichever cloning method was the choice, the
476 inserted gene piece would be included in the primers, which would be designed, synthesized, and used in the

477 cloning. This step, leading to a further modified Spike with the furin-cleavage site added at the S1/S2 junction,
478 could be completed in no more than two weeks.

479 Unlike the engineering of Spike, no complicated design is needed here, except that the RdRp gene segment
480 from RaBtCoV/4991 would need to be included. Gibson assembly could have been used here. In this technique,
481 several fragments, each adjacent pair sharing 20-40 bp overlap, are combined together in one simple reaction
482 to assemble a long DNA product. Two or three fragments, each covering a significant section of the ORF1b
483 gene, would be selected based on known bat coronavirus sequences. One of these fragments would be the RdRp
484 segment of RaBtCoV/4991 83 . Each fragment would be PCR amplified with proper overlap regions introduced
485 in the primers. Finally, all purified fragments would be pooled in equimolar concentrations and added to the
486 Gibson reaction mixture, which, after a short incubation, would yield the desired ORF1b gene in whole.

487 Step 4: Produce the designed viral genome using reverse genetics and recover live viruses (0.5 month)

488 Reverse genetics have been frequently used in assembling whole viral genomes, including coronavirus genomes
489 67,[92][93][94][95][96] . The most recent example is the reconstruction of the SARS-CoV-2 genome using the
490 transformation-assisted recombination in yeast 97 . Using this method, the Swiss group assembled the entire
491 viral genome and produced live viruses in just one week 97 . This efficient technique, which would not leave any
492 trace of artificial manipulation in the created viral genome, has been available since 2017 98,99 . In addition to
493 the engineered spike gene (from steps 1 and 2) and the ORF1b gene (from step 3), other fragments covering the
494 rest of the genome would be obtained either through RT-PCR amplification from the template virus or through
495 DNA synthesis by following a sequence slightly altered from that of the template virus. We believe that the
496 latter approach was more likely as it would allow sequence changes introduced into the variable regions of less
497 conserved proteins, the process of which could be easily guided by multiple sequence alignments. The amino
498 acid sequences of more conserved functions, such as that of the E protein, might have been left unchanged.
499 All DNA fragments would then be pooled together and transformed into yeast, where the cDNA version of the
500 SARS-CoV-2 genome would be assembled via transformation-assisted recombination. Of course, an alternative
501 method of reverse genetics, one of which the WIV has successfully used in the past 67 , could also be employed
502 67,[92][93][94][95][96]100 . Although some earlier reverse genetics approaches may leave restriction sites at where
503 different fragments would be joined, these traces would be hard to detect as the exact site of ligation can be
504 anywhere in the ~30kb genome. Either way, a cDNA version of the viral genome would be obtained from the
505 reverse genetics experiment. Subsequently, *in vitro* transcription using the cDNA as the template would yield
506 the viral RNA genome, which upon transfection into Vero E6 cells would allow the production of live viruses
507 bearing all of the designed properties.

508 Step 5: Optimize the virus for fitness and improve its hACE2-binding affinity *in vivo* (2.5-3 months) Virus
509 recovered from step 4 needs to be further adapted undergoing the classic experiment -serial passage in laboratory
510 animals 101 . This final step would validate the virus' fitness and ensure its receptor oriented adaptation toward
511 its intended host, which, according to the analyses above, should be human. Importantly, the RBM and the
512 furin-cleavage site, which were introduced into the Spike protein separately, would now be optimized together
513 as one functional unit. Among various available animal models (e.g. mice, hamsters, ferrets, and monkeys) for
514 coronaviruses, hACE2 transgenic mice (hACE2-mice) should be the most proper and convenient choice here.
515 This animal model has been established during the study of SARS-CoV and has been available in the Jackson
516 Laboratory for many years [102][103][104] .

517 The procedure of serial passage is straightforward. Briefly, the selected viral strain from step 4, a precursor
518 of SARS-CoV-2, would be intranasally inoculated into a group of anaesthetized hACE2-mice. Around 2-3 days
519 post infection, the virus in lungs would usually amplify to a peak titer. The mice would then be sacrificed and
520 the lungs homogenized. Usually, the mouse-lung supernatant, which carries the highest viral load, would be
521 used to extract the candidate virus for the next round of passage. After approximately 10-15 rounds of passage,
522 the hACE2-binding affinity, the infection efficiency, and the lethality of the viral strain would be sufficiently
523 enhanced and the viral genome stabilized 101 . Finally, after a series of characterization experiments (e.g. viral
524 kinetics assay, antibodies response assay, symptom observation and pathology examination), the final product,
525 SARS-CoV-2, would be obtained, concluding the whole creation process. From this point on, this viral pathogen
526 could be amplified (most probably using Vero E6 cells) and produced routinely.

527 It is noteworthy that, based on the work done on SARS-CoV, the hACE2-mice, although suitable for SARS-
528 CoV-2 adaptation, is not a good model to reflect the virus' transmissibility and associated clinical symptoms in
529 humans. We believe that those scientists might not have used a proper animal model (such as the golden Syrian
530 hamster) for testing the transmissibility of SARS-CoV-2 before the outbreak of COVID-19. If they had done this
531 experiment with a proper animal model, the highly contagious nature of SARS-CoV-2 would be extremely evident
532 and consequently SARS-CoV-2 would not have been described as "not causing human-tohuman transmission" at
533 the start of the outbreak.

534 We also speculate that the extensive laboratory adaptation, which is oriented toward enhanced transmissibility
535 and lethality, may have driven the virus too far. As a result, SARS-CoV-2 might have lost the capacity to
536 attenuate on both transmissibility and lethality during its current adaptation in the human population. This
537 hypothesis is consistent with the lack of apparent attenuation of SARS-CoV-2 so far despite its great prevalence
538 and with the observation that a recently emerged, predominant variant only shows improved transmissibility
539 [105][106][107][108] .

Serial passage is a quick and intensive process, where the adaptation of the virus is accelerated. Although intended to mimic natural evolution, serial passage is much more limited in both time and scale. As a result, less random mutations would be expected in serial passage than in natural evolution. This is particularly true for conserved viral proteins, such as the E protein. Critical in viral replication, the E protein is a determinant of virulence and engineering of it may render SARS-CoV-2 attenuated [109][110][111]. Therefore, at the initial assembly stage, these scientists might have decided to keep the amino acid sequence of the E protein unchanged from that of ZC45/ZXC21. Due to the conserved nature of the E protein and the limitations of serial passage, no amino acid mutation actually occurred, resulting in a 100% sequence identity on the E protein between SARS-CoV-2 and ZC45/ZXC21. The same could have happened to the marks of molecular cloning (restriction sites flanking the RBM). Serial passage, which should have partially naturalized the SARS-CoV-2 genome, might not have removed all signs of artificial manipulation.

Many questions remain unanswered about the origin of SARS-CoV-2. Prominent virologists have implicated in a Nature Medicine letter that laboratory escape, while not being entirely ruled out, was unlikely and that no sign of genetic manipulation is present in the SARS-CoV-2 genome [4]. However, here we show that genetic evidence within the spike gene of SARS-CoV-2 genome (restriction sites flanking the RBM; tandem rare codons used at the inserted furin-cleavage site) does exist and suggests that the SARS-CoV-2 genome should be a product of genetic manipulation. Furthermore, the proven concepts, well-established techniques, and knowledge and expertise are all in place for the convenient creation of this novel coronavirus in a short period of time.


Motives aside, the following facts about SARS-CoV-2 are well-supported:

1. If it was a laboratory product, the most critical element in its creation, the backbone/template virus (ZC45/ZXC21), is owned by military research laboratories. 2. The genome sequence of SARS-CoV-2 has likely undergone genetic engineering, through which the virus has gained the ability to target humans with enhanced virulence and infectivity. 3. The characteristics and pathogenic effects of SARS-CoV-2 are unprecedented. The virus is highly transmissible, onset-hidden, multi-organ targeting, sequelae-unclear, lethal, and associated with various symptoms and complications. 4. SARS-CoV-2 caused a world-wide pandemic, taking millions of lives and shutting down the global economy. It has a destructive power like no other.

Judging from the evidence that we and others have gathered, we believe that finding the origin of SARS-CoV-2 should involve an independent audit of the WIV P4 laboratories and the laboratories of their close collaborators. Such an investigation should have taken place long ago and should not be delayed any further.

We also note that in the publication of the chimeric virus SHC015-MA15 in 2015, the attribution of funding of Zhengli Shi by the NIAID was initially left out. It was reinstated in the publication in 2016 in a corrigendum, perhaps after the meeting in January 2016 to reinstate NIH funding for gain-of-function research on viruses. This is an unusual scientific behavior, which needs an explanation for.

What is not thoroughly described in this report is the various evidence indicating that several coronaviruses recently published (RaTG13 [18], RmYN02 [30], and several pangolin coronaviruses [27][28][29][31]) are highly suspicious and likely fraudulent. These fabrications would serve no purpose other than to deceive the scientific community and the general public so that the true identity of SARS-CoV-2 is hidden. Although exclusion of details of such evidence does not alter the conclusion of the current report, we do believe that these details would provide additional support for our contention that SARS-CoV-2 is a laboratory-enhanced virus and a product of gain-of-function research. A follow-up report focusing on such additional evidence is now being prepared and will be submitted shortly.



2

Figure 1: Figure 2 :




Figure 2: Figure 3 :




Figure 3: Figure 4 :




Figure 4: Figure 5 :




Figure 5: Figure 6 :

Human SARS-CoV BJ01	655 - GICASYHTVSL- - - - -RSTS - 670
Human SARS-CoV CUHK-W1	655 - GICASYHTVSL- - - - -RSTS - 670
Human SARS-CoV Tor2	655 - GICASYHTVSL- - - - -RSTS - 670
Human SARS-CoV Frankfurt-1	655 - GICASYHTVSL- - - - -RSTS - 670
Human SARS-CoV Urbani	655 - GICASYHTVSL- - - - -RSTS - 670
Civet SARS-CoV civet020	655 - GICASYHTVSSL- - - - -RSTS - 670
Civet SARS-CoV sz16	655 - GICASYHTVSSL- - - - -RSTS - 670
Racoon dog SARS-CoV A030	655 - GICASYHTVSSL- - - - -RSTS - 670
SARS-CoV-2	669 - GICASYQTQTNSPRRARSVA - 688
Pangolin CoV MP789	n/a - GICASYQTQTNS- - - - -RSVS - n/a
Bat SARSr-CoV RaTG13	669 - GICASYQTQTNS- - - - -RSVA - 684
Bat SARSr-CoV LYRa11	659 - GICASYHTASLL- - - - -RNTD - 674
Bat SARSr-CoV LYRa3	659 - GICASYHTASLL- - - - -RNTG - 674
Bat SARSr-CoV RsSHC014	656 - GICASYHTVSSL- - - - -RSTS - 671
Bat SARSr-CoV Rs4084	656 - GICASYHTVSSL- - - - -RSTS - 671
Bat SARSr-CoV WIV1	656 - GICASYHTVSSL- - - - -RSTS - 671
Bat SARSr-CoV Rs3367	656 - GICASYHTVSSL- - - - -RSTS - 671
Bat SARSr-CoV Rs7327	656 - GICASYHTVSSL- - - - -RSTS - 671
Bat SARSr-CoV Rs9401	656 - GICASYHTVSSL- - - - -RSTS - 671
Bat SARSr-CoV Rs4231	655 - GICASYHTVSSL- - - - -RSTS - 670
Bat SARSr-CoV WIV16	655 - GICASYHTVSSL- - - - -RSTS - 670
Bat SARSr-CoV Rs4874	655 - GICASYHTVSSL- - - - -RSTS - 670
Bat SARSr-CoV ZC45	646 - GICASYHTASIL- - - - -RSTS - 661
Bat SARSr-CoV ZXC21	645 - GICASYHTASIL- - - - -RSTG - 660
Bat SARSr-CoV Rf4092	634 - GICASYHTASTL- - - - -RGVG - 649
Bat SARSr-CoV Rf/JL2012	636 - GICASYHTASLL- - - - -RSTG - 651
Bat SARSr-CoV JTMCI5	636 - GICASYHTASLL- - - - -RSTG - 651
Bat SARSr-CoV 16BO133	636 - GICASYHTASLL- - - - -RSTG - 651
Bat SARSr-CoV B15-21	636 - GICASYHTASLL- - - - -RSTG - 651
Bat SARSr-CoV YN2013	633 - GICASYHTASTL- - - - -RSIG - 648
Bat SARSr-CoV Anlong-103	633 - GICASYHTASTL- - - - -RSVG - 648
Bat SARSr-CoV Rp/Shaanxi2011	640 - GICASYHTASVL- - - - -RSTG - 655
Bat SARSr-CoV Rs/HuB2013	641 - GICASYHTASVL- - - - -RSTG - 656
Bat SARSr-CoV YNLF/34C	641 - GICASYHTASVL- - - - -RSTG - 656
Bat SARSr-CoV YNLF/31C	641 - GICASYHTASVL- - - - -RSTG - 656
Bat SARSr-CoV Rf1	641 - GICASYHTASHL- - - - -RSTG - 656
Bat SARSr-CoV 273	641 - GICASYHTASHL- - - - -RSTG - 656
Bat SARSr-CoV Rf/SX2013	639 - GICASYHTASLL- - - - -RSTG - 654
Bat SARSr-CoV Rf/HeB2013	641 - GICASYHTASLL- - - - -RSTG - 656
Bat SARSr-CoV Cp/Yunnan2011	641 - GICASYHTASLL- - - - -RNTG - 656
Bat SARSr-CoV Rs672	641 - GICASYHTASTL- - - - -RSVG - 656
Bat SARSr-CoV Rs4255	641 - GICASYHTASTL- - - - -RSVG - 656
Bat SARSr-CoV 4081	641 - GICASYHTASTL- - - - -RSVG - 656
Bat SARSr-CoV Rm1	641 - GICASYHTASVL- - - - -RSTG - 656
Bat SARSr-CoV 279	641 - GICASYHTASVL- - - - -RSTG - 656
Bat SARSr-CoV Rs/GX2013	642 - GICASYHTASVL- - - - -RSTG - 657
Bat SARSr-CoV Rs806	641 - GICASYHTASLL- - - - -RSTG - 656
Bat SARSr-CoV HKU3-1	642 - GICASYHTASVL- - - - -RSTG - 657
Bat SARSr-CoV Longquan-140	642 - GICASYHTASVL- - - - -RSTG - 657
Bat SARSr-CoV Rp3	641 - GICASYHTASTL- - - - -RSVG - 656
Bat SARSr-CoV Rs4247	642 - GICASYHTASTL- - - - -RSVG - 657
Bat SARSr-CoV Rs4237	641 - GICASYHTASTL- - - - -RSVG - 656
Bat SARSr-CoV As6526	641 - GICASYHTASTL- - - - -RSVG - 656

Fault

tat cag act cag act aat tct **cct** **cgg** **cgg** **gca** cgt agt gta gct agt caa tcc atc att
Y Q T Q T N S **P** **R** **R** **A** R S V A S Q S I I

Figure 7:

1 Acknowledgements

We would like to thank Daoyu Zhang for sharing with us the findings of mutations in the E proteins in different sub-groups of ? coronaviruses. We also thank all the anonymous scientists and other individuals, who have contributed in uncovering various facts associated with the origin of SARS-CoV-2.

2 II. Remarks

- [China Honors Ian Lipkin ()] , <https://www.publichealth.columbia.edu/public-health-now/news/china-honorsian-lipkin> *China Honors Ian Lipkin* 2020.
- [Holmes and Academic ()] , E Holmes , C V Academic . <https://www.sydney.edu.au/AcademicProfiles/profile/resource?urlid=edward.holmes&type=cv> 2020.
- [Sequencing and Preprint ()] , Provenance Sequencing , Preprint . <https://zenodo.org/record/3987503#.Xz9GzC-z3GI> 2020.
- [Agnihothram ()] *A mouse model for Betacoronavirus subgroup 2c using a bat coronavirus strain HKU5 variant*, S Agnihothram . 2014. 5 p. .
- [Roberts ()] *A mouse-adapted SARS coronavirus causes disease and mortality in BALB/c mice*, A Roberts . 2007. 3 p. e5.
- [Hoffmann et al. ()] ‘A Multibasic Cleavage Site in the Spike Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells’. M Hoffmann , H Kleine-Weber , S Pohlmann . *Mol* 2020. 78 p. e5.
- [Wu ()] *A new coronavirus associated with human respiratory disease in China*, F Wu . 2020. 579 p. .
- [Zhou ()] ‘A Novel Bat Coronavirus Closely Related to SARS-CoV-2 Contains Natural Insertions at the S1/S2 Cleavage Site of the Spike Protein’. H Zhou . *Curr* 2020. 30 p. e3.
- [Zhou ()] ‘A pneumonia outbreak associated with a new coronavirus of probable bat origin’. P Zhou . *Nature* 2020.
- [Menachery ()] ‘A SARS-like cluster of circulating bat coronaviruses shows potential for human emergence’. V D Menachery . *Nat* 2015. 21 p. .
- [Belouzard et al. ()] ‘Activation of the SARS coronavirus spike protein via sequential proteolytic cleavage at two distinct sites’. S Belouzard , V C Chu , G R Whittaker . *Proc Natl Acad Sci* 2009. p. .
- [Bengston ()] *All journal articles evaluating the origin or epidemiology of SARS-CoV-2 that utilize the RaTG13 bat strain genomics are potentially flawed and should be retracted*, D Bengston . 10.31219/osf.io/wy89d. 2020. (OSF Preprints)
- [Xie ()] ‘An Infectious cDNA Clone of SARS-CoV-2’. X Xie . *Cell Host* 2020. 27 p. e3.
- [Hou ()] ‘Angiotensin-converting enzyme 2 (ACE2) proteins of different bat species confer variable susceptibility to SARS-CoV entry’. Y Hou . *Arch* 2010. 155 p. .
- [Roberts ()] ‘Animal models and vaccines for SARS-CoV infection’. A Roberts . *Virus* 2008. 133 p. .
- [Zhang] *Anomalies in BatCoV/RaTG13*, D Zhang .
- [Robinson ()] *Another expert challenges assertions that SARS-CoV-2 was not genetically engineered*, C Robinson . <https://gmwatch.org/en/news/latest-news/> 19383, 2020.
- [Kam ()] *Antibodies against trimeric S glycoprotein protect hamsters against SARS-CoV challenge despite their capacity to mediate FcγRII-dependent entry into B cells in vitro*, Y W Kam . 2007. 25 p. .
- [Liu ()] *Are pangolins the intermediate host of the 2019 novel coronavirus (SARS-CoV-2)?*, P Liu . 2020. 16 p. e1008421.
- [Lau ()] ‘Attenuated SARS-CoV-2 variants with deletions at the S1/S2 junction’. S Y Lau . *Emerg Microbes* 2020. 9 p. .
- [Muth ()] ‘Attenuation of replication by a 29 nucleotide deletion in SARS-coronavirus acquired during the early stages of human-to-human transmission’. D Muth . *Sci* 2018. 8 p. 15177.
- [Zeng ()] ‘Bat Severe Acute Respiratory Syndrome-Like Coronavirus WIV1 Encodes an Extra Accessory Protein, ORFX, Involved in Modulation of the Host Immune Response’. L P Zeng . *J* 2016. 90 p. .
- [Shang ()] ‘Cell entry mechanisms of SARS-CoV-2’. J Shang . *Proc Natl Acad Sci* 2020. p. .
- [Vashee ()] *Cloning, Assembly, and Modification of the Primary Human Cytomegalovirus Isolate Toledo by Yeast-Based Transformation-Associated Recombination*, S Vashee . 2017. p. e2.
- [Ge ()] ‘Coexistence of multiple coronaviruses in several bat colonies in an abandoned mineshaft’. X Y Ge . *Virol* 2016. 31 p. .
- [Almazan ()] ‘Construction of a severe acute respiratory syndrome coronavirus infectious cDNA clone and a replicon to study coronavirus RNA synthesis’. F Almazan . *J* 2006. 80 p. .

- 634 [Schoeman and Fielding ()] ‘Coronavirus envelope protein: current knowledge’. D Schoeman , B C Fielding .
635 *Virol* 2019. 16 p. 69.
- 636 [Song et al. ()] *Cryo-EM structure of the SARS coronavirus spike glycoprotein in complex with its host cell*
637 *receptor ACE2*, W Song , M Gui , X Wang , Y Xiang . 2018. 14 p. e1007236.
- 638 [Singla et al. ()] *Denovo Assembly of RaTG13 Genome Reveals Inconsistencies Further Obscuring SARS-CoV-2*
639 *Origins*, M Singla , S Ahmad , C Gupta , T Sethi . doi: 10.20944/ preprints202008.0595.v1. 2020. Preprints.
640 p. 2020080595.
- 641 [Ge ()] ‘Detection of alpha-and beta coronaviruses in rodents from Yunnan’. X Y Ge . *China. Virol* 2017. 14 p.
642 98.
- 643 [Ren ()] ‘Difference in receptor usage between severe acute respiratory syndrome (SARS) coronavirus and SARS-
644 like coronavirus of bat origin’. W Ren . *J* 2008. 82 p. .
- 645 [Hu ()] *Discovery of a rich gene pool of bat SARS-related coronaviruses provides new insights into the origin of*
646 *SARS coronavirus*, B Hu . 2017. 13 p. e1006698.
- 647 [Luo ()] ‘Discovery of Novel Bat Coronaviruses in South China That Use the Same Receptor as Middle East
648 Respiratory Syndrome Coronavirus’. C M Luo . *J* 2018. 92.
- 649 [Li ()] ‘Emergence of SARS-CoV-2 through Recombination and Strong Purifying Selection’. X Li . *bioRxiv* 2020.
- 650 [Almazan et al. ()] ‘Engineering infectious cDNAs of coronavirus as bacterial artificial chromosomes’. F Almazan
651 , S Marquez-Jurado , A Nogales , L Enjuanes . *Methods Mol Biol* 2015. p. .
- 652 [Watanabe ()] ‘Entry from the cell surface of severe acute respiratory syndrome coronavirus with cleaved S
653 protein as revealed by pseudotype virus bearing cleaved S protein’. R Watanabe . *J* 2008. 82 p. .
- 654 [Ren ()] ‘Full-length genome sequences of two SARS-like coronaviruses in horseshoe bats and genetic variation
655 analysis’. W Ren . *J Gen* 2006. 87 p. .
- 656 [Follis et al. ()] *Furin cleavage of the SARS coronavirus spike glycoprotein enhances cell-cell fusion but does not*
657 *affect virion entry*, K E Follis , J York , J H Nunberg . 2006. 350 p. .
- 658 [Wu et al. ()] *Furin, a potential therapeutic target for COVID-19*, Canrong Wu , YY , Yang Liu , Peng
659 Zhang , Yali Wang , Hua Li , Qiqi Wang , Yang Xu , Mingxue Li , Mengzhu Zheng , Lixia Chen .
660 <http://www.chinaxiv.org/abs/202002.00062> 2020. (Preprint (chinaXiv))
- 661 [Ito ()] ‘Generation of a highly pathogenic avian influenza A virus from an avirulent field isolate by passaging in
662 chickens’. T Ito . *J* 2001. 75 p. .
- 663 [Ommeh ()] ‘Genetic Evidence of Middle East Respiratory Syndrome Coronavirus (MERS-Cov) and Widespread
664 Seroprevalence among Camels in Kenya’. S Ommeh . *Virol* 2018. 33 p. .
- 665 [Oldfield ()] ‘Genome-wide engineering of an infectious clone of herpes simplex virus type 1 using synthetic
666 genomics assembly methods’. L M Oldfield . *Proc Natl Acad Sci* 2017. p. .
- 667 [Hu ()] ‘Genomic characterization and infectivity of a novel SARS-like coronavirus in Chinese bats’. D Hu .
668 *Emerg Microbes* 2018. 7 p. 154.
- 669 [Drexler ()] ‘Genomic characterization of severe acute respiratory syndrome-related coronavirus in European
670 bats and classification of coronaviruses based on partial RNA-dependent RNA polymerase gene sequences’. J
671 F Drexler . *J* 2010. 84 p. .
- 672 [Wang ()] *hACE2 Transgenic Mouse Model For Coronavirus (COVID-19) Research. The Jackson Labora-*
673 *tory Research Highlight*, Q Wang . [https://www.jax.org/news-and-insights/2020/february/](https://www.jax.org/news-and-insights/2020/february/introducingmouse-model-for-corona-virus#)
674 [introducingmouse-model-for-corona-virus#](https://www.jax.org/news-and-insights/2020/february/introducingmouse-model-for-corona-virus#) 2020.
- 675 [Claas ()] *Human influenza A H5N1 virus related to a highly pathogenic avian influenza virus*, E C Claas . 1998.
676 351 p. .
- 677 [Liu ()] ‘Identification of common deletions in the spike protein of SARS-CoV-2’. Z Liu . *J Virol* 2020.
- 678 [Lam ()] ‘Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins’. T T Lam . *Nature* 2020.
- 679 [Takayama ()] ‘In Vitro and Animal Models for SARS-CoV-2 research’. K Takayama . *Trends Pharmacol* 2020.
680 41 p. 513517.
- 681 [Yuan ()] ‘Intraspecies diversity of SARS-like coronaviruses in *Rhinolophus sinicus* and its implications for the
682 origin of SARS coronaviruses in humans’. J Yuan . *J Gen* 2010. 91 p. .
- 683 [Segreto and Deigin ()] *Is considering a geneticmanipulation origin for SARS-CoV-2 a conspiracy theory*
684 *that must be censored?*, R Segreto , Y Deigin . DOI: 10.13140/ RG.2.2.31358.13 129/1. 2020. (Preprint
685 (Researchgate))
- 686 [Ge ()] *Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor*, X Y Ge .
687 2013. 503 p. .

- 688 [Yang ()] ‘Isolation and Characterization of a Novel Bat Coronavirus Closely Related to the Direct Progenitor
689 of Severe Acute Respiratory Syndrome Coronavirus’. X L Yang . *J* 2015. 90 p. .
- 690 [Guan ()] *Isolation and characterization of viruses related to the SARS coronavirus from animals in southern
691 China*, Y Guan . 2003. 302 p. .
- 692 [Xiao ()] ‘Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins’. K Xiao . *Nature* 2020.
- 693 [Robinson ()] *Journals censor lab origin theory for SARS-CoV-2*, C Robinson . [https://www.gmwatch.org/
694 en/news/latest-news/19475-journals-censor-lab-origin-theory-for-sars-cov-2](https://www.gmwatch.org/en/news/latest-news/19475-journals-censor-lab-origin-theory-for-sars-cov-2) 2020.
- 695 [Luo ()] ‘Longitudinal Surveillance of Betacoronaviruses in Fruit Bats in Yunnan Province, China During’. Y
696 Luo . *Viol Sin33*, 2009-2016. 2018. p. .
- 697 [Lin and Chen ()] *Major Concerns on the Identification of Bat Coronavirus Strain RaTG13 and Quality of
698 Related Nature Paper*, X Lin , S Chen . 2020. Preprints. p. 2020060044.
- 699 [Wu et al. ()] ‘Mechanisms of host receptor adaptation by severe acute respiratory syndrome coronavirus’. K Wu
700 , G Peng , M Wilken , R J Geraghty , F Li . *J Biol* 2012. 287 p. .
- 701 [Yeung ()] ‘MERS coronavirus induces apoptosis in kidney and lung by upregulating Smad7 and FGF2’. M L
702 Yeung . *Nat* 2016. 1 p. 16004.
- 703 [Chu ()] ‘MERS coronaviruses from camels in Africa exhibit region-dependent genetic diversity’. D K W Chu .
704 *Proc Natl Acad Sci* 2018. p. .
- 705 [Zhou et al. ()] *Middle East respiratory syndrome coronavirus infection: virus-host cell interactions and
706 implications on pathogenesis*, J Zhou , H Chu , J F Chan , K Y Yuen . 2015. 12 p. 218.
- 707 [Chan ()] ‘Middle East respiratory syndrome coronavirus: another zoonotic betacoronavirus causing SARS-like
708 disease’. J F Chan . *Clin Microbiol* 2015. 28 p. .
- 709 [Sirotkin and Sirotkin ()] ‘Might SARS-CoV-2 Have Arisen via Serial Passage through an Animal Host or
710 Cell Culture?’. K Sirotkin , D Sirotkin . 10.1002/bies.202000091. [https://doi.org/10.1002/bies.
711 202000091](https://doi.org/10.1002/bies.202000091) *BioEssays* 2020.
- 712 [Sun et al. ()] ‘Modifications to the hemagglutinin cleavage site control the virulence of a neurotropic H1N1
713 influenza virus’. X Sun , L V Tse , A D Ferguson , G R Whittaker . *J* 2010. 84 p. .
- 714 [Wan ()] ‘Molecular Mechanism for Antibody-Dependent Enhancement of Coronavirus Entry’. Y Wan . *J* 2020.
715 94.
- 716 [Mou ()] ‘Mutations from bat ACE2 orthologs markedly enhance ACE2-Fc neutralization of SARSCoV-2’. H
717 Mou . 10.1101/2020.06.29.178459. <https://doi.org/10.1101/2020.06.29.178459> *BioRxiv* 2020.
- 718 [Lee ()] ‘No evidence of coronaviruses or other potentially zoonotic viruses in Sunda pangolins (*Manis javanica*)
719 entering the wildlife trade via Malaysia’. J Lee . 10.1101/2020.06.19.158717. [https://doi.org/10.1101/
720 2020.06.19.158717](https://doi.org/10.1101/2020.06.19.158717) *bioRxiv* 2020.
- 721 [Maiti ()] *On The Origin of SARS-CoV-2 Virus*, A K Maiti . 10.22541/au.159355977.76503625. 2020. (Preprint
722 (authorea.com))
- 723 [Cui et al. ()] ‘Origin and evolution of pathogenic coronaviruses’. J Cui , F Li , Z L Shi . *Nat Rev* 2019. 17 p.
724 181192.
- 725 [Sia ()] ‘Pathogenesis and transmission of SARS-CoV-2 in golden hamsters’. S F Sia . *Nature* 2020.
- 726 [Piplani et al. ()] S Piplani , P K Singh , D A Winkler , N Petrovsky . arXiv:2005.06199. *silico comparison of
727 spike protein-ACE2 binding affinities across species; significance for the possible origin of the SARS-CoV-2
728 virus. arXiv*, 2020.
- 729 [Wang ()] *Preliminary investigation of viruses carried by bats on the southeast coastal area (?????
730 ????????????)*, Y Wang . 2017. (Master Thesis)
- 731 [Zhang et al. ()] ‘Probable Pangolin Origin of SARS-CoV-2 Associated with the COVID19 Outbreak’. T Zhang
732 , Q Wu , Z Zhang . *Curr* 2020. 30 p. 1578.
- 733 [Seyran ()] ‘Questions concerning the proximal origin of SARS-CoV-2’. M Seyran . *J Med Virol* 2020.
- 734 [Thao ()] ‘Rapid reconstruction of SARS-CoV-2 using a synthetic genomics platform’. T T N Thao . *Nature* 2020.
- 735 [Li ()] ‘Receptor recognition mechanisms of coronaviruses: a decade of structural studies’. F Li . *J Virol* 89 2015.
736 p. 195464.
- 737 [Poon ()] ‘Recurrent mutations associated with isolation and passage of SARS coronavirus in cells from non-
738 human primates’. L L Poon . *J Med* 2005. 76 p. .
- 739 [Kuo et al. ()] ‘Retargeting of coronavirus by substitution of the spike glycoprotein ectodomain: crossing the
740 host cell species barrier’. L Kuo , G J Godeke , M J Raamsman , P S Masters , P J Rottier . *J* 2000. 74 p.
741 1393406.

- 742 [Yount ()] ‘Reverse genetics with a full-length infectious cDNA of severe acute respiratory syndrome coronavirus’.
743 B Yount . *Proc Natl Acad Sci* 2003. p. .
- 744 [Scobey ()] ‘Reverse genetics with a full-length infectious cDNA of the Middle East respiratory syndrome
745 coronavirus’ . T Scobey . *Proc Natl Acad Sci* 2013. p. .
- 746 [Kido ()] ‘Role of host cellular proteases in the pathogenesis of influenza and influenza-induced multiple organ
747 failure’ . H Kido . *BiochimBiophys Acta1824* 2012. p. .
- 748 [Zhang ()] *SARS-CoV-2 Could Come from a Lab -A Critique of*, B Zhang . https://www.linkedin.com/pulse/sars-cov-2-could-come-from-labcritique-proximal-origin-billy-zhang?articleId=6651628681431175168#comments6651628681431175168&trk=public_profile_article_view 2020. (The Proximal Origin of SARS-CoV-2. Published in Nature Medicine.)
- 752 [Zhan et al. ()] ‘SARS-CoV-2 is well adapted for humans. What does this mean for re-emergence?’. S H Zhan ,
753 B E Deverman , Y A Chan . 10.1101/2020.05.01.073262. *BioRxiv* 2020.
- 754 [Menachery ()] ‘SARS-like WIV1-CoV poised for human emergence’ . V D Menachery . *Proc Natl Acad Sci* 2016.
755 p. .
- 756 [Scientific evidence and logic behind the claim that the Wuhan coronavirus is man-made ()] *Scientific evidence
757 and logic behind the claim that the Wuhan coronavirus is man-made*, <https://nerdhaspower.weebly.com>
758 2020.
- 759 [Nieto-Torres ()] *Severe acute respiratory syndrome coronavirus envelope protein ion channel activity promotes
760 virus fitness and pathogenesis*, J L Nieto-Torres . 2014. 10 p. e1004077.
- 761 [Lau ()] ‘Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats’ . S K Lau . *Proc
762 Natl Acad Sci* 2005. p. .
- 763 [Chan and Zhan ()] ‘Single source of pangolin CoVs with a near identical Spike RBD to SARSCoV-2’ . Y A Chan
764 , S H Zhan . 10.1101/2020.07.07.184. *bioRxiv* 2020. 374.
- 765 [Plante ()] ‘Spike mutation D614G alters SARS-CoV-2 fitness and neutralization susceptibility’ . J A Plante .
766 10.1101/2020.09.01.278689. *bioRxiv* 2020.
- 767 [Yurkovetskiy ()] ‘Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant’ . L
768 Yurkovetskiy . 10.1101/2020.07.04.187757. *bioRxiv* 2020.
- 769 [Shang ()] ‘Structural basis of receptor recognition by SARS-CoV-2’ . J Shang . *Nature* 2020.
- 770 [Pervushin ()] *Structure and inhibition of the SARS coronavirus envelope protein ion channel*, K Pervushin .
771 2009. 5 p. e1000511.
- 772 [Li et al. ()] *Structure of SARS coronavirus spike receptor-binding domain complexed with receptor*, F Li , W Li
773 , M Farzan , S C Harrison . 2005. 309 p. .
- 774 [Li ()] ‘Structure, Function, and Evolution of Coronavirus Spike Proteins’ . F Li . *Annu Rev* 2016. 3 p. .
- 775 [Becker ()] ‘Synthetic recombinant bat SARS-like coronavirus is infectious in cultured cells and in mice’ . M M
776 Becker . *Proc Natl Acad Sci* 2008. p. .
- 777 [Yount et al. ()] ‘Systematic assembly of a full-length infectious cDNA of mouse hepatitis virus strain A59’ . B
778 Yount , M R Denison , S R Weiss , R S Baric . *J* 2002. 76 p. .
- 779 [Rahalkar and bahulikar ()] ‘The Abnormal Nature of the Fecal Swab Sample used for NGS Analysis of RaTG13
780 Genome Sequence Imposes a Question on the Correctness of the RaTG13 Sequence’ . M Rahalkar , R
781 &bahulikar . *Preprints.org* 2020. p. 2020080205.
- 782 [Zhang ()] ‘The D614G mutation in the SARS-CoV-2 spike protein reduces S1 shedding and increases infectivity’ .
783 L Zhang . 10.1101/2020.06.12.148726. <https://doi.org/10.1101/2020.06.12.148726> *bioRxiv* 2020.
- 784 [Sørensen et al. ()] *The Evidence which Suggests that This Is No Naturally Evolved Virus*, B
785 Sørensen , A Dalglish , A & susrud . [https://www.minervanett.no/files/2020/07/13/](https://www.minervanett.no/files/2020/07/13/TheEvidenceNoNaturalEvol.pdf)
786 [TheEvidenceNoNaturalEvol.pdf](https://www.minervanett.no/files/2020/07/13/TheEvidenceNoNaturalEvol.pdf) 2020. (Preprint)
- 787 [Zhang ()] ‘The ORF8 Protein of SARS-CoV-2 Mediates Immune Evasion through Potently Downregulating
788 MHC-I’ . Y Zhang . 10.1101/2020.05.24.111823. <https://doi.org/10.1101/2020.05.24.111823>
789 *bioRxiv* 2020.
- 790 [Zhang ()] *The Pan-SL-CoV/GD sequences may be from contamination*, D Zhang . 10.5281/zenodo.3885333.
791 2020. (Preprint (zenodo.org))
- 792 [Andersen et al. ()] ‘The proximal origin of SARSCoV-2’ . K G Andersen , A Rambaut , W I Lipkin , E C Holmes
793 , R F Garry . *Nat* 2020. 26 p. .
- 794 [Cheng ()] *The S2 Subunit of QX-type Infectious Bronchitis Coronavirus Spike Protein Is an Essential
795 Determinant of Neurotropism*, J Cheng . 2019. p. 11.

- 796 [Hassanin ()] ‘The SARS-CoV-2-like virus found in captive pangolins from Guangdong should be better
797 sequenced’. A Hassanin . <https://doi.org/10.1101/2020.05.07.077016>. *bioRxiv* 2020.
- 798 [Coutard ()] ‘The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent
799 in CoV of the same clade’. B Coutard . *Antiviral* 2020. 176 p. 104742.
- 800 [Korber ()] *Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19*
801 *Virus*, B Korber . 2020. 182 p. e19.
- 802 [Yang ()] ‘Two Mutations Were Critical for Batto-Human Transmission of Middle East Respiratory Syndrome
803 Coronavirus’. Y Yang . *J* 2015. 89 p. .
- 804 [Rahalkar and bahulikar ()] *Understanding the Origin of ‘BatCoVraTG13’, a Virus Closest to SARS-CoV-2.*
805 *Preprints*, M C Rahalkar , R A &bahulikar . 2020. p. 2020050322.
- 806 [Johnson et al. ()] *Viral metagenomics, protein structure, and reverse genetics: Key strategies for investigating*
807 *coronaviruses*, B A Johnson , R L Graham , V D &menachery . 2018. 517 p. .
- 808 [Robinson ()] *Was the COVID-19 virus genetically engineered*, C Robinson . [https://gmwatch.org/en/](https://gmwatch.org/en/news/latestnews/)
809 [news/latestnews/](https://gmwatch.org/en/news/latestnews/) 19383, 2020.